



# CLIP

15 minute Worksheet



Helping the patient with reduced hydration and nutrition

## 7: Managing a gastrostomy

Advanced level

Produced by  
**Coleman Education  
 Centre**  
**St. Oswald's Hospice**  
 Regent Avenue, Gosforth  
 Newcastle-upon-Tyne  
 NE3 1EE  
 Tel: 0191 285 0063  
 Fax: 0191 284 8004

This version written and edited by:  
**Dorothy Matthews**  
 Senior Nurse  
**Lynn Gibson**  
 Senior physiotherapist  
 Northgate Hospital Learning Disability  
 Palliative Care Team

**Claud Regnard**  
 Consultant in Palliative Medicine  
 St. Oswald's Hospice, Newcastle  
 Hospitals NHS Trust and  
 Northgate&Prudhoe NHS Trust  
**Margaret Kindlen**, Head of Education  
 St. Oswald's Hospice

Development of this worksheet was  
 supported by  
**the NHS Executive  
 BEACON initiative**

Northgate Palliative Care Team  
 NHS Beacon site

### Aim of this worksheet

To understand the principles of managing a gastrostomy.

### How to use this worksheet

- You can work through this worksheet by yourself, or with a tutor.
- Read the case study below, then work on the questions overleaf.
- The work page is on the right side, the information page is on the left.
- Work any way you want: you can try answering from your own knowledge (in which case fold over the information page), you can use the information page (this is not cheating- you learn as you find the information), or you can use other sources of information
- It should take you about 15 minutes. If anything is unclear, discuss it with a colleague.
- If you think any information is wrong or out of date let us know
- Use the activity on the back page and take this learning into your workplace.

### Case study

**Ben is a 33 year old man who has a moderate learning disability together with hydrocephalus spastic diplegia, visual impairment and epilepsy. He has been diagnosed as having a carcinoma of kidney with lung metastases. A decline in Ben's nutritional status meant decisions relating to maintaining his weight and general overall condition had to be made. As Ben has a relatively good quality of life it was decided by Ben's family and the palliative care team to arrange for Ben to have a percutaneous endoscopic gastrostomy (PEG). The staff caring for Ben would like further advice on aftercare and problems relating to the PEG.**

## Gastrostomy

- T. It is an opening through the abdominal wall into the stomach through which a feeding tube can be passed.
  - F. PEG = percutaneous endoscopic gastrostomy since an endoscope is used to insert the tube.
  - F. X-ray control or surgery can be used (only surgery needs a general anaesthetic).
  - T. With good after care, gastrostomies can last for years (see also worksheet on *Non-oral routes*).
  - T. In a patient with a swallowing disorder, the risk is higher with the a nasogastric tube.
- Initially Ben was given a PEG with a small tube exiting from the stoma site. This was changed after six weeks to a low profile PEG tube 'button' which is discreet and comfortable to use and is favoured especially by young people.

### Initial care of the PEG site (if you have a local PEG team, follow their protocols)

#### *For a new gastrostomy*

1. The external fixation plate should be left intact for 72 hours to minimise gastric leakage and allow adherence of the stoma to abdominal wall.
2. Several days after the procedure, the external fixation plate should be moved slightly to prevent adherence.
3. Each day:
  - always wash hands before caring for PEG sites.
  - clean the newly formed stoma site using sterile saline solution.
  - rotate the tube to avoid adherence to the tract.

#### *When the stoma site has healed:*

- Cleaning should be done daily using water or mild soap solution.
- A dressing is not necessary unless there is a heavy discharge.
- Bathing and showering are allowed at this stage.
- The tube should be flushed with 25-50 mls of water between feeds and after drug administration to prevent blockages. The frequency of flushes and the amount of water used will depend on frequency of feeds and whether medication is being given through the PEG.

## Possible problems

### Redness around the stoma:

*Skin Excoriation:* Is the fixation too tight or not being cleaned regularly? Is it too loose causing leakage? This inflammation and reddened skin can be prevented by use of barrier creams. If the stoma site is leaking gastric contents, ensure that the fixation plate is in the correct position.

*Local infection:* Are you using good hygiene techniques? Infection occurs in 3-15% of cases so a good hygiene technique is essential. Dressings may lead to moisture accumulation. Turning the tube daily prevents crusts forming and the development of scar tissue. If a local cellulitis is suspected (redness spreading well beyond the stoma) antibiotics should be given through the gastrostomy (usually flucloxacillin) and a wound swab should be taken for culture and sensitivity.

**Tube blockage:** why has it happened? Can it be reduced by regular flushing? Try flushing with a small diameter syringe to exert more pressure. If water does not clear the blockage, Soda water or carbonised liquid may clear the tube. Pineapple juice contains a proteolytic enzyme which can be used to digest a blockage caused by feeds. Regular use of cranberry juice can reduce blockages. Some drugs coagulate with feeds and cause blockage (eg. anacids, carbamazepine, phenytoin) and they need to be well diluted (at least an equal volume of water) or given at least one hour separate from a feed.

**Dislodged tube:** has this been pulled too hard? If this is a new gastrostomy of 10 days or less contact the team who inserted the tube. If the gastrostomy is well established then a urinary Foley catheter should be inserted in the first instant to keep the stoma patent until a replacement can be obtained.

**Nausea and vomiting** this may due to the infusion running too fast or volumes being too large. Some patients need a prokinetic (metoclopramide or domperidone) to encourage gastric emptying.

**Chest infections:** these can be due to aspiration of feeds. There is no evidence that non-oral feeding reduces the risk of aspiration. Maintain head of bed at minimum of 30 degrees may help and consider using metoclopramide to increase gastric emptying.

### After care

- Oral hygiene should not be overlooked. Problems can be avoided with good regular oral hygiene as a part of routine care.
- Ben should be weighed regularly, his weight recorded, intake recorded and a check made that the prescribed amount is given. All of these are less important once his cancer progresses.
- Ben's tube size and date of insertion and removal method should be recorded.
- Details for delivery of supplies and feed for Ben should be made available.
- Balloon inflated tubes should be regularly checked (sizes varies 5-30ml).
- Be innovative with volumes and schedules. NB. A flexible feeding regime constitutes a good quality of life.
- Stratton (2002) found that 83% of patients on home artificial nutrition still had the desire to eat.

# MICQ

Write down your answers to these questions:

- A gastrostomy passes food directly to the stomach      True      False
- PEG stands for Percutaneous Exterior Gastrostomy      True      False
- Insertion needs a general anaesthetic      True      False
- A gastrostomy can be used long-term      True      False
- The risk of aspiration is lower with a gastrostomy      True      False

## Write

What can you do for the following problems?

Problem	Your solution
Redness around the stoma	
Blocked tube	
Dislodged tube	
Vomiting after feeds	
Increase in chest infections	

## Reflect

Think about what after-care Ben will need

## FURTHER ACTIVITY: Managing a gastrostomy

- The next time you have a patient with a gastrostomy:
  - list the advantages for that patient
  - observe any problems the patient is having and consider ways of helping

## FURTHER READING : Managing a gastrostomy

### Journal articles

- Arrowsmith H. Nursing Management of patients receiving gastrostomy feeding. *British Journal of Nursing* 1996; **5**: 268-73.
- Boyd K.J. Beeken L. Tube feeding in palliative care: benefits and problems *Palliative Medicine*, 1994; **8**: 156-158
- Campos ACL, Butters M and Meguid MM. Home enteral nutrition via gastrostomy in advanced head and neck cancer patients. *Head and Neck* 1990; **12**: 137-42.
- Dwolatzky T. Berezovski S. Friedmann R. Paz J. Clarfield AM. Stessman J. Hamburger R. Jaul E. Friedlander Y. Rosin A. Sonnenblick M. A prospective comparison of the use of nasogastric and percutaneous endoscopic gastrostomy tubes for long-term enteral feeding in older people. *Clinical Nutrition*. 2001; **20**(6): 535-40.
- Gauderer MW. Percutaneous endoscopic gastrostomy and the evolution of contemporary long-term enteral access. *Clinical Nutrition*. 2002; **21**(2):103-10.
- Goodhall L. Tube feeding dilemmas: can artificial nutrition and hydration be legally or ethically withheld or withdrawn? *Journal of Advanced Nursing* 1997, **25**: 217-222
- Howell M. Do nurses know enough about percutaneous endoscopic gastrostomy?. *Nursing Times*. 2002; **98**(17): 40-2.
- Kimber CP, Beasley SW. Limitations of percutaneous endoscopic gastrostomy in facilitating enteral nutrition in children: review of the shortcomings of a new technique. *Journal of Paediatric and Child Health*, 1999; **35**: 427-31.
- Laing B, Smithers M and Harper J. Percutaneous fluoroscopic gastrostomy: a safe option? *Medical Journal of Australia* 1994; **161**: 308-10.
- Larson D. Burton DD. Scgroeder KW. Percutaneous endoscopic Gastrostomy. Indicates success, complications and mortality in 314 consecutive patients. *Gastroenterology* 1987; **93**: 48-52.
- Mandal A, Steel A, Davidson AR, Ashby C. Day-case percutaneous endoscopic gastrostomy: a viable proposition? *Postgraduate Medical Journal*, 2000; **76**: 157-9.
- Myssiorek D, Siegel D, Vambutas A. Fluoroscopically placed gastrostomies in the head and neck patient. *Laryngoscope*, 1998; **108**: 1557-60.
- Norton B, Homer-Ward M, Donnelly MT, Long RG, Holmes GKT. A randomised prospective comparison of percutaneous endoscopic gastrostomy and nasogastric tube feeding after dysphagic stroke. *British Medical Journal*, 1996; **312**: 13-6.
- Pennington C. To PEG or not to PEG. *Clinical Medicine*. 2002; **2**(3): 250-5.
- Raha S. Woodhouse K. Who should have a PEG? *Age and Ageing* 1993; **22**: 313-315.
- Stratton R. Use and benefits of enteral nutrition. *Primary Health Care*, 2002; **12**(2): 25-30.
- White S. Percutaneous endoscopic Gastrostomy P.E.G. *Nursing Standard* 1998; **12**: April 1-7.

# CLIP

**Current Learning In Palliative care**  
An accessible learning programme for health care professionals

### Fifty seven 15 minute worksheets are available on:

- An introduction to palliative care (3 worksheets)
- Helping the patient with pain (9 worksheets)
- Helping the patient with symptoms other than pain (11 worksheets)
- Moving the ill patient (2 worksheets)
- Psychological needs (8 worksheets)
- Helping patients with reduced hydration and nutrition (8 worksheets)
- Procedures in palliative care (4 worksheets)
- Understanding and helping the person with alternative communication (learning disabilities) (5 worksheets)
- The last hours and days (4 worksheets)
- Bereavement (3 worksheets)

© 2004

Helping the Patient with Advanced Disease: a Workbook  
Regnard C. ed.

Oxford: Radcliffe Medical Press [www.radcliffe-oxford.com](http://www.radcliffe-oxford.com)